

ORAL PRESENTATION

Open Access

The determinants of *C. difficile* infection in long-term care facilities: a portrait of patient- and facility-level factors across 90 care regions in the veterans affairs health care system

KA Brown^{1*}, M Jones¹, F Adler², M Leecaster¹, K Nechodom³, V Stevens⁴, M Samore¹, J Mayer¹

From 3rd International Conference on Prevention and Infection Control (ICPIC 2015)
Geneva, Switzerland. 16-19 June 2015

Introduction

Clostridium difficile infection (CDI) is an infectious diarrheal disease that is associated with antibiotic and healthcare exposures. Although individual-level risk factors have been extensively studied, the facility-level factors that drive CDI have not.

Objectives

To study the determinants of CDI incidence across long term care (LTC) facilities, with a specific interest in the role importation of infectious patients from acute care (AC) facilities.

Methods

We conducted a retrospective cohort study of CDI from 2006 through 2012 across Veterans Affairs local health-care systems (HCS) where both AC and LTC patient censuses were above an average of 10 patients per day. Our outcome was LTC-onset *C. difficile* lab-identified event, defined as a case with onset ≥ 3 days after admission occurring at least 8 weeks from a previous positive test.

Results

We identified 90 local HCS that met our inclusion criteria. The incidence of *C. difficile* infection in LTC facilities was 3.6 per 10,000 patient-days. In bivariate weighted linear regression analyses, the most important predictors of facility CDI incidence were importation ($R^2=0.63$, $p<0.001$) and antibiotic prescribing ($R^2=0.58$,

$p<0.001$). Time-series analyses revealed that increases in *C. difficile* case importation from AC facilities preceded increases in CDI rates for a period of up to 8 weeks. Multi-level analyses, that included individual-level covariates, revealed that *C. difficile* importation and facility-level antibiotic use acted independently of resident age, direct antibiotic exposure and direct proton pump inhibitor use.

Conclusion

This is the first study showing that importation of *C. difficile* cases from AC facilities and facility-level antibiotic use are principal drivers of CDI in LTC facilities. A regional approach, addressing rates in AC facilities, is needed to control CDI in LTC facilities.

Disclosure of interest

None declared.

Authors' details

¹Epidemiology, University of Utah, Salt Lake City, UT, USA. ²Mathematics, University of Utah, Salt Lake City, UT, USA. ³University of Utah, Salt Lake City, UT, USA. ⁴Pharmacy, University of Utah, Salt Lake City, UT, USA.

Published: 16 June 2015

doi:10.1186/2047-2994-4-S1-O36

Cite this article as: Brown et al.: The determinants of *C. difficile* infection in long-term care facilities: a portrait of patient- and facility-level factors across 90 care regions in the veterans affairs health care system. *Antimicrobial Resistance and Infection Control* 2015 **4**(Suppl 1):O36.

¹Epidemiology, University of Utah, Salt Lake City, UT, USA
Full list of author information is available at the end of the article